Complete Weight Enumerator of a Family of Linear Codes from Cyclotomy

Shudi Yang and Zheng-An Yao and Chang-An Zhao

arXiv:1507.05732v1 [cs.IT] 21 Jul 2015

Abstract

Linear codes have been an interesting topic in both theory and practice for many years. In this paper, for a prime p, we determine the explicit complete weight enumerators of a family of linear codes over $\mathbb{F}_p$ with defining set related to cyclotomy. These codes may have applications in cryptography and secret sharing schemes.

Index Terms

Linear code, complete weight enumerator, cyclotomy, Gaussian period, period polynomial.

I. INTRODUCTION

Throughout this paper, let p be a prime and $r = p^m$ for some positive integer m. Denote by $\mathbb{F}_p$ a finite field with p elements. An $[n, \kappa, \delta]$ linear code C over $\mathbb{F}_p$ is a κ-dimensional subspace of $\mathbb{F}_p^n$ with minimum distance δ [1], [2].

The complete weight enumerator [1], [3] of a code C over $\mathbb{F}_p$ enumerates the codewords according to the number of symbols of each kind contained in each codeword. Denote the field elements by $\mathbb{F}_p = \{w_0, w_1, \cdots, w_{p-1}\}$, where $w_0 = 0$. Also let $\mathbb{F}_p^*$ denote $\mathbb{F}_p \backslash \{0\}$. For a codeword $\mathsf{c} = (c_0, c_1, \cdots, c_{n-1}) \in \mathbb{F}_p^n$, let $w[\mathsf{c}]$ be the complete weight enumerator of c defined as

$$w[\mathsf{c}] = w_0^{k_0} w_1^{k_1} \cdots w_{p-1}^{k_{p-1}},$$

where k_j is the number of components of c equal to w_j, $\sum_{j=0}^{p-1} k_j = n$. The complete weight enumerator of the code C is then

$$\mathrm{CWE}(C) = \sum_{\mathsf{c} \in C} w[\mathsf{c}].$$

The weight distribution of a linear code has been studied extensively for decades and we refer the reader to [4], [5], [6], [7], [8], [9], [10], [11], [12], [13], [14], [15], [16] and references therein for an overview of the related researches. It is not difficult to see that the complete weight enumerators are just the (ordinary) weight enumerators for binary linear codes. For nonbinary linear codes, the weight enumerators can be obtained from their complete weight enumerators.

The information of the complete weight enumerator of a linear code is of vital use in theories and practical applications. For instance, Blake and Kith investigated the complete weight enumerator of Reed-Solomon codes and showed that they could be helpful in soft decision decoding [3], [17]. In [18], the study of the monomial and quadratic bent functions was related to the complete weight enumerators of linear codes. It was illustrated by Ding $et\ al.$ [19], [20] that the complete weight enumerator can be applied to calculate the deception probabilities of certain authentication codes. In [21], [22], [23], the authors studied the complete weight enumerators of some constant composition codes and presented some families of optimal constant composition codes.

However, it is generally an extremely difficult problem to evaluate the complete weight enumerator of linear codes and there are few information on this topic in literature besides the above mentioned [3], [17], [21], [22], [23]. Kuzmin and Nechaev considered the generalized Kerdock code and related linear codes over Galois rings and determined their complete weight enumerators in [24] and [25]. Very recently, the complete weight enumerators of linear codes over finite fields were studied in [26], [27], [28], [29], [30].

In [2], [31], [32], [33], [34], Ding $et\ al.$ proposed a generic method to construct linear codes with a few nonzero weights by employing trace function. We introduce this construction below.

Let $D = \{d_1, d_2, \cdots, d_z\} \subseteq \mathbb{F}_r$ for a positive integer z. Denote by Tr the trace function from $\mathbb{F}_r$ to $\mathbb{F}_p$. A linear code of length z over $\mathbb{F}_p$ is defined by

$$C_D = \{(\mathrm{Tr}(xd))_{d \in D} : x \in \mathbb{F}_r\}, \tag{1}$$

and D is called the defining set of this code C_D.

S.D. Yang is with the Department of Mathematics, Sun Yat-sen University, Guangzhou 510275, and School of Mathematical Sciences, Qufu Normal University, Shandong 273165, P.R.China.

Z.-A. Yao and C.-A Zhao are with the Department of Mathematics, Sun Yat-sen University, Guangzhou 510275, P.R.China.

E-mail: yangshd3@mail2.sysu.edu.cn, mcsyao@mail.sysu.edu.cn

Manuscript received *********; revised ********.

The authors in [2], [31], [32], [33], [34] presented such linear codes and investigated their weight enumerators for some well chosen defining sets. Along this research line, the authors of [27] and [29] investigated the complete weight enumerators of linear codes with defining sets for certain special cases.

Let α be a fixed primitive element of $\mathbb{F}_r$ and $r-1=nN$ for positive integers $n>1$ and $N>1$. We always assume that $N|\frac{r-1}{p-1}$. The cyclotomic classes of order N are defined by $C_i^{(N,r)}=\alpha^i\left\langle\alpha^N\right\rangle$ for $i=0,1,\cdots,N-1$, where $\langle\alpha^N\rangle$ denotes the subgroup of $\mathbb{F}_r^*$ generated by α^N.

In [31], Ding and Niederreiter constructed two classes of cyclotomic linear codes $C_{\bar{D}}$ over $\mathbb{F}_p$ of order 3 with defining sets $\bar{D}=D_0$ and $\bar{D}=D_0\bigcup D_1$, where $D_j=\{\alpha^j\alpha^{3i}:i=0,1,\cdots,\frac{r-1}{3(p-1)}-1\}$ for $j=0,1$. Inspired by the original idea proposed in [31], we shall study the complete weight enumerator of C_D with defining sets constructed from cyclotomy.

In this paper, the defining set D of the code C_D is given by

$$D=\bigcup_{i\in I}C_i^{(N,r)}, \tag{2}$$

where $I\subset\{0,1,\cdots,N-1\}$ with $\#I=l>0$.

Therefore the code C_D defined by (1) with defining set D of (2) is a linear code with length nl and dimension at most m. Obviously, the different choices of I lead to different codes C_D.

We employ Gaussian periods to determine the complete weight enumerator of C_D. A general formula is given for the defining set D of (2). Moreover, as applications of this formula, we explicitly present the complete weight enumerator of C_D for the special cases of $N=3$ and $N=4$, respectively. In fact, the defining set of [31] is a complete set of coset representatives of the factor group $C_0^{(3,r)}/\mathbb{F}_p^*$. We generalize it to the whole coset $C_i^{(3,r)}$ with $i=0,1,2$. As it turns out that, the codes C_D for $N=3$ and $N=4$ are linear codes with few weights. More precisely, they have nonzero weights not more than $four$ and thus will have many applications in cryptography [35] and secret sharing schemes [36]. They can be employed to construct constant composition codes [21], [37] which have important use in communications engineering [38]. We also give some examples to illustrate our results, which shows that some of these codes are optimal or have the best parameters due to Ding's tables [2].

It should be remarked that, when $D=C_0^{(N,r)}$, our result is as same as that of [26]. We give the result by Gaussian periods and the authors of [26] gave the result by Gauss sums. They are equivalent though in different manner. In addition, when $N=3$, since $C_0^{(3,r)}=\{xy:x\in\mathbb{F}_p^* \ \text{ and } \ y\in D_0\}=(\mathbb{F}_p^*)D_0$, the weight enumerator of $C_{\bar{D}}$ in [31] can be obtained from that of C_D. Thus, we extend the results in [31] to some extent. These will be shown in details in the consequent sections.

The remainder of this paper is organized as follows. Section II recalls some definitions and results about cyclotomic classes and Gaussian periods which will be useful in the sequel. Section III presents the complete weight enumerator of the code C_D with defining set D, including a general strategy and the special case of $N=3$ and $N=4$. Section III concludes this paper.

II. Mathematical foundation of Cyclotomy in $\mathbb{F}_r$

In this section, we introduce some necessary mathematical foundation which will be of use in the sequel.

Recall that p is a prime, $r=p^m$ and $r-1=nN$ for two positive integers $n>1$ and $N>1$. Let α be a fixed primitive element of $\mathbb{F}_r$. Define $C_i^{(N,r)}=\alpha^i\left\langle\alpha^N\right\rangle$ for $i=0,1,\cdots,N-1$, where $\langle\alpha^N\rangle$ denotes the subgroup of $\mathbb{F}_r^*$ generated by α^N. The cosets $C_i^{(N,r)}$ are called the *cyclotomic classes* of order N. It is easily seen that $\#C_i^{(N,r)}=\frac{r-1}{N}=n$ and $C_i^{(N,r)}=C_{i \pmod N}^{(N,r)}$. Moreover, $C_0^{(N,r)},\ C_1^{(N,r)},\cdots,C_{N-1}^{(N,r)}$ and $\{0\}$ form a partition of $\mathbb{F}_r$.

The *Gaussian periods* are defined by

$$\eta_i^{(N,r)}=\sum_{x\in C_i^{(N,r)}}\zeta_p^{\mathrm{Tr}(x)},$$

where $\zeta_p=e^{2\pi\sqrt{-1}/p}$ and Tr is the trace function from $\mathbb{F}_r$ to $\mathbb{F}_p$. Clearly $\eta_i^{(N,r)}=\eta_{i \pmod N}^{(N,r)}$ if $i\geqslant N$.

Generally speaking, it is very hard to compute the values of Gaussian periods. They are known only in a few cases and they can be obtained from *period polynomials* $\Psi_{(N,r)}(X)$ which are defined by

$$\Psi_{(N,r)}(X)=\prod_{i=0}^{N-1}(X-\eta_i^{(N,r)}).$$

It was shown that $\Psi_{(N,r)}(X)$ is a polynomial with integer coefficients [39]. The following two lemmas, which were cited from [39], will be of use in the sequel.

Lemma 1. [39] *Let $N=3$ and $r=p^m$. We have the following results on the factorization of $\Psi_{(3,r)}(X)$.*
(a) If $p\equiv 2\pmod 3$, then $m\equiv 0\pmod 2$, and

$$\Psi_{(3,r)}(X)=\begin{cases}3^{-3}(3X+1+2\sqrt{r})(3X+1-\sqrt{r})^2 & \text{if } \ m\equiv 0 \pmod 4,\\ 3^{-3}(3X+1-2\sqrt{r})(3X+1+\sqrt{r})^2 & \text{if } \ m\equiv 2 \pmod 4.\end{cases}$$

(b) If $p \equiv 1 \pmod 3$*, and* $m \equiv 0 \pmod 3$*, then*

$$\Psi_{(3,r)}(X) = 3^{-3}(3X+1-s_1r^{\frac{1}{3}})(3X+1+\frac{1}{2}(s_1+9t_1)r^{\frac{1}{3}})(3X+1+\frac{1}{2}(s_1-9t_1)r^{\frac{1}{3}}).$$

where s_1 *and* t_1 *are given by* $4r^{\frac{1}{3}} = s_1^2 + 27t_1^2$, $s_1 \equiv 1 \pmod 3$ *and* $\gcd(s_1, p) = 1$.

Lemma 2. [39] *Let* $N = 4$ *and* $r = p^m$*. We have the following results on the factorization of* $\Psi_{(4,r)}(X)$*.*
(a) If $p \equiv 3 \pmod 4$*, then* $m \equiv 0 \pmod 2$*, and*

$$\Psi_{(4,r)}(X) = \begin{cases} 4^{-4}(4X+1+3\sqrt{r})(4X+1-\sqrt{r})^3 & \text{if} \quad m \equiv 0 \pmod 4, \\ 4^{-4}(4X+1-3\sqrt{r})(4X+1+\sqrt{r})^3 & \text{if} \quad m \equiv 2 \pmod 4. \end{cases}$$

(b) If $p \equiv 1 \pmod 4$*, and* $m \equiv 0 \pmod 4$*, then*

$$\begin{aligned} \Psi_{(4,r)}(X) &= 4^{-4}\left((4X+1)+\sqrt{r}+2r^{\frac{1}{4}}u_1\right)\left((4X+1)+\sqrt{r}-2r^{\frac{1}{4}}u_1\right) \\ &\quad \times \left((4X+1)-\sqrt{r}+4r^{\frac{1}{4}}v_1\right)\left((4X+1)-\sqrt{r}-4r^{\frac{1}{4}}v_1\right), \end{aligned}$$

where u_1 *and* v_1 *are given by* $r^{\frac{1}{2}} = u_1^2 + 4v_1^2$, $u_1 \equiv 1 \pmod 4$ *and* $\gcd(u_1, p) = 1$.

III. MAIN RESULTS

We maintain al notations from the previous sections, and we want now to determine the complete weight enumerator of the codes C_D defined by (1) with defining set D of (2). These codes may have different property with different defining set D. Thus, we focus on a general case and two special cases.

A. A general strategy

Recall that

$$C_D = \{(\mathrm{Tr}(xd))_{d\in D} : x \in \mathbb{F}_r\},$$

where $D = \bigcup_{i\in I} C_i^{(N,r)}$ with $I \subset \{0, 1, \cdots, N-1\}$ and $\#I = l > 0$.

Clearly $x = 0$ gives the zero codeword which contributes w_0^{nl} to the complete weight enumerator. Thus, we only need to focus on $x \in \mathbb{F}_r^*$.

Let

$$N_k(\rho) = \#\{d \in D : \mathrm{Tr}(xd) = \rho \text{ and } \ x \in C_k^{(N,r)}\},$$

where $0 \leqslant k \leqslant N-1$.

Note that $N|\frac{r-1}{p-1}$ leads to $\mathbb{F}_p^* \subseteq C_0^{(N,r)}$. Thus, we can deduce that

$$\begin{aligned} N_k(\rho) &= \sum_{d\in D}\frac{1}{p}\sum_{y\in\mathbb{F}_p}\zeta_p^{y(\mathrm{Tr}(xd)-\rho)} \\ &= \frac{nl}{p} + \frac{1}{p}\sum_{y\in\mathbb{F}_p^*}\zeta_p^{-y\rho}\sum_{d\in D}\zeta_p^{y\mathrm{Tr}(xd)} \\ &= \frac{nl}{p} + \frac{1}{p}\sum_{y\in\mathbb{F}_p^*}\zeta_p^{-y\rho}\left(\sum_{i\in I}\sum_{d\in C_i^{(N,r)}}\zeta_p^{\mathrm{Tr}(yxd)}\right) \\ &= \frac{nl}{p} + \frac{1}{p}\sum_{y\in\mathbb{F}_p^*}\zeta_p^{-y\rho}\sum_{i\in I}\eta_{k+i}^{(N,r)} \\ &= \begin{cases} \frac{nl}{p} + \frac{p-1}{p}\sum_{i\in I}\eta_{k+i}^{(N,r)} & \text{if} \quad \rho = 0, \\ \frac{nl}{p} - \frac{1}{p}\sum_{i\in I}\eta_{k+i}^{(N,r)} & \text{otherwise}, \end{cases} \end{aligned} \tag{3}$$

where the fourth equality holds since $\mathbb{F}_p^* \subseteq C_0^{(N,r)}$.

Therefore, it follows from Equation (3) that

$$CWE(C_D) = w_0^{nl} + n\sum_{k=0}^{N-1} w_0^{\frac{nl}{p}+\frac{p-1}{p}\sum_{i\in I}\eta_{k+i}^{(N,r)}}\prod_{\rho=1}^{p-1} w_\rho^{\frac{nl}{p}-\frac{1}{p}\sum_{i\in I}\eta_{k+i}^{(N,r)}}, \tag{4}$$

where $n = \#C_k^{(N,r)} = \frac{r-1}{N}$.

B. Linear codes from cyclotomy of order N=3

In this subsection, we will give the complete weight enumerator of the code C_D with defining set D constructed from cyclotomy of order $N=3$.

The results on Gaussian periods of order $N=3$ follow immediately from Lemma 1.

Lemma 3. *Let $N=3$, $r=p^m$ and $N|\frac{r-1}{p-1}$. We have the following results on Gaussian periods.*
(a) If $p\equiv 1 \pmod 3$, then $m\equiv 0 \pmod 3$ and

$$\begin{aligned}\eta_0^{(3,r)} &= -\frac{1-s_1r^{\frac{1}{3}}}{3},\\ \eta_1^{(3,r)} &= -\frac{2+(s_1+9t_1)r^{\frac{1}{3}}}{6},\\ \eta_2^{(3,r)} &= -\frac{2+(s_1-9t_1)r^{\frac{1}{3}}}{6},\end{aligned}$$

where s_1 and t_1 are defined in Lemma 1.
(b) If $p\equiv 2 \pmod 3$, then $m\equiv 0 \pmod 2$.
For the case of $m\equiv 0 \pmod 4$, we have

$$\begin{aligned}\eta_0^{(3,r)} &= -\frac{1+2\sqrt{r}}{3},\\ \eta_1^{(3,r)} &= \eta_2^{(3,r)}=-\frac{1-\sqrt{r}}{3}.\end{aligned}$$

For the case of $m\equiv 2 \pmod 4$, we have

$$\begin{aligned}\eta_0^{(3,r)} &= -\frac{1-2\sqrt{r}}{3},\\ \eta_1^{(3,r)} &= \eta_2^{(3,r)}=-\frac{1+\sqrt{r}}{3}.\end{aligned}$$

The following gives the complete weight enumerator of the code C_D with defining set D constructed from cyclotomy of order $N=3$.

Theorem 4. *Let $N=3$, $r=p^m$ and $N|\frac{r-1}{p-1}$. Let $\#I=2$ and $r-1=nN$. Then the code C_D of* (1) *is a $[2n,m]$ linear code over $\mathbb{F}_p$ and its complete weight enumerator is given as follows.*
(a) If $p\equiv 1 \pmod 3$ and $m\equiv 0 \pmod 3$, then

$$\begin{aligned}CWE(C_D) &= w_0^{2n}+nw_0^{\frac{2n}{p}-\frac{p-1}{6p}(4-(s_1-9t_1)r^{\frac{1}{3}})}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{1}{6p}(4-(s_1-9t_1)r^{\frac{1}{3}})}\\ &\quad +nw_0^{\frac{2n}{p}-\frac{p-1}{3p}(2+s_1r^{\frac{1}{3}})}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{1}{3p}(2+s_1r^{\frac{1}{3}})}\\ &\quad +nw_0^{\frac{2n}{p}-\frac{p-1}{6p}(4-(s_1+9t_1)r^{\frac{1}{3}})}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{1}{6p}(4-(s_1+9t_1)r^{\frac{1}{3}})}.\end{aligned}$$

(b) If $p\equiv 2 \pmod 3$ and $m\equiv 0 \pmod 4$, then

$$\begin{aligned}CWE(C_D) &= w_0^{2n}+2nw_0^{\frac{2n}{p}-\frac{p-1}{3p}(2+\sqrt{r})}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{1}{3p}(2+\sqrt{r})}\\ &\quad +nw_0^{\frac{2n}{p}-\frac{2(p-1)}{3p}(1-\sqrt{r})}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{2}{3p}(1-\sqrt{r})}.\end{aligned}$$

(c) If $p \equiv 2 \pmod 3$ *and* $m \equiv 2 \pmod 4$*, then*

$$\begin{aligned} CWE(C_D) &= w_0^{2n} + 2nw_0^{\frac{2n}{p}-\frac{p-1}{3p}(2-\sqrt{r})}\prod_{\rho=1}^{p-1} w_\rho^{\frac{2n}{p}+\frac{1}{3p}(2-\sqrt{r})} \\ &\quad + nw_0^{\frac{2n}{p}-\frac{2(p-1)}{3p}(1+\sqrt{r})}\prod_{\rho=1}^{p-1} w_\rho^{\frac{2n}{p}+\frac{2}{3p}(1+\sqrt{r})}. \end{aligned}$$

Proof: It follows from Lemma 3 that the following assertions hold.
(a) If $p \equiv 1 \pmod 3$ and $m \equiv 0 \pmod 3$, then

$$\begin{aligned} \eta_0^{(3,r)} + \eta_1^{(3,r)} &= -\frac{4-(s_1-9t_1)r^{\frac{1}{3}}}{6}, \\ \eta_1^{(3,r)} + \eta_2^{(3,r)} &= -\frac{2+s_1r^{\frac{1}{3}}}{3}, \\ \eta_2^{(3,r)} + \eta_0^{(3,r)} &= -\frac{4-(s_1+9t_1)r^{\frac{1}{3}}}{6}, \end{aligned}$$

where s_1 and t_1 are defined in Lemma 1.
(b) If $p \equiv 2 \pmod 3$ and $m \equiv 0 \pmod 4$, then

$$\begin{aligned} \eta_0^{(3,r)} + \eta_1^{(3,r)} &= -\frac{2+\sqrt{r}}{3} = \eta_2^{(3,r)} + \eta_0^{(3,r)}, \\ \eta_1^{(3,r)} + \eta_2^{(3,r)} &= -\frac{2(1-\sqrt{r})}{3}. \end{aligned}$$

(c) If $p \equiv 2 \pmod 3$ and $m \equiv 2 \pmod 4$, then

$$\begin{aligned} \eta_0^{(3,r)} + \eta_1^{(3,r)} &= -\frac{2-\sqrt{r}}{3} = \eta_2^{(3,r)} + \eta_0^{(3,r)}, \\ \eta_1^{(3,r)} + \eta_2^{(3,r)} &= -\frac{2(1+\sqrt{r})}{3}. \end{aligned}$$

The desired conclusions then follow from Equation (4). ■

When $\#I = 1$, the result is straightforward from Lemma 3 and Equation (4).

Theorem 5. *Let* $N = 3$*,* $r = p^m$ *and* $N|\frac{r-1}{p-1}$*. Let* $\#I = 1$ *and* $r - 1 = nN$*. Then the code* C_D *of* (1) *is an* $[n, m]$ *cyclic code over* $\mathbb{F}_p$ *and its complete weight enumerator is given as follows.*
(a) If $p \equiv 1 \pmod 3$ *and* $m \equiv 0 \pmod 3$*, then*

$$\begin{aligned} CWE(C_D) &= w_0^{n} + nw_0^{\frac{n}{p}-\frac{p-1}{3p}(1-s_1r^{\frac{1}{3}})}\prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{3p}(1-s_1r^{\frac{1}{3}})} \\ &\quad + nw_0^{\frac{n}{p}-\frac{p-1}{6p}(2+(s_1+9t_1)r^{\frac{1}{3}})}\prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{6p}(2+(s_1+9t_1)r^{\frac{1}{3}})} \\ &\quad + nw_0^{\frac{n}{p}-\frac{p-1}{6p}(2+(s_1-9t_1)r^{\frac{1}{3}})}\prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{6p}(2+(s_1-9t_1)r^{\frac{1}{3}})}, \end{aligned}$$

where s_1 *and* t_1 *are defined in Lemma 1.*
(b) If $p \equiv 2 \pmod 3$ *and* $m \equiv 0 \pmod 4$*, then*

$$\begin{aligned} CWE(C_D) &= w_0^{n} + 2nw_0^{\frac{n}{p}-\frac{p-1}{3p}(1-\sqrt{r})}\prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{3p}(1-\sqrt{r})} \\ &\quad + nw_0^{\frac{n}{p}-\frac{p-1}{3p}(1+2\sqrt{r})}\prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{3p}(1+2\sqrt{r})}. \end{aligned}$$

(c) If $p \equiv 2 \pmod 3$ *and* $m \equiv 2 \pmod 4$*, then*

$$\begin{aligned} CWE(C_D) &= w_0^n + 2nw_0^{\frac{n}{p}-\frac{p-1}{3p}(1+\sqrt{r})}\prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{3p}(1+\sqrt{r})} \\ &\quad +nw_0^{\frac{n}{p}-\frac{p-1}{3p}(1-2\sqrt{r})}\prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{3p}(1-2\sqrt{r})}. \end{aligned}$$

Remark 6. It should be remarked that the authors in [31] determined the weight enumerators of two classes of cyclotomic linear codes $C_{\bar{D}}$ over $\mathbb{F}_p$ of order 3 with defining sets $\bar{D} = D_0$ and $\bar{D} = D_0 \bigcup D_1$, where $D_j = \{\alpha^j\alpha^{3i} : i = 0, 1, \cdots, \frac{r-1}{3(p-1)} - 1\}$ for $j = 0, 1$.

Note that $C_0^{(3,r)} = \{xy : x \in \mathbb{F}_p^* \ \text{ and } \ y \in D_0\} = (\mathbb{F}_p^*)D_0$. By Theorems 4 and 5, we can obtain the weight enumerator of C_D for $D = \bigcup_{i\in I} C_i^{(3,r)}$ with $I \subset \{0, 1, 2\}$. Dividing each nonzero weight of C_D with $p-1$ yields the weight enumerator of $C_{\bar{D}}$, which conforms to the results described in [31]. Thus we generalize the results of [31] to some extent.

Example 7. Let $(p, m) = (2, 6)$ and $N = 3$. Then $r = 64$ and $n = 21$. Suppose that α is a primitive element of $\mathbb{F}_{64}$.

(1) For the case of $D = C_1^{(3,r)} = \alpha\langle\alpha^3\rangle$.

By Theorem 5, the code C_D of (1) is a binary $[21, 6, 8]$ cyclic code and its complete weight enumerator is

$$CWE(C_D) = w_0^{21} + 21w_0^{13}w_1^8 + 42w_0^9w_1^{12},$$

which is consistent with numerical computation by Magma.

This code is the best binary cyclic code of length 21 and dimension 6 according to the tables given by Ding [2].

(2) For the case of $D = C_0^{(3,r)} \bigcup C_1^{(3,r)} = \langle\alpha^3\rangle \bigcup \alpha\langle\alpha^3\rangle$.

By Theorem 4, the code C_D of (1) is a binary $[42, 6, 20]$ cyclic code which is optimal with respect to the Griesmer bound, and its complete weight enumerator is

$$CWE(C_D) = w_0^{42} + 42w_0^{22}w_1^{20} + 21w_0^{18}w_1^{24},$$

which is consistent with numerical computation by Magma.

C. Linear codes from cyclotomy of order N=4

In this subsection, we will present the complete weight enumerator of the code C_D with defining set D constructed from cyclotomy of order $N = 4$.

The results on Gaussian periods of order $N = 4$ follow immediately from Lemma 2.

Lemma 8. *Let* $N = 4$, $r = p^m$ *and* $N|\frac{r-1}{p-1}$*. We have the following results on Gaussian periods.*
(a) If $p \equiv 1 \pmod 4$*, then* $m \equiv 0 \pmod 4$ *and*

$$\begin{aligned} \eta_0^{(4,r)} &= -\frac{1+\sqrt{r}+2r^{\frac{1}{4}}u_1}{4}, \\ \eta_1^{(4,r)} &= -\frac{1-\sqrt{r}+4r^{\frac{1}{4}}v_1}{4}, \\ \eta_2^{(4,r)} &= -\frac{1+\sqrt{r}-2r^{\frac{1}{4}}u_1}{4}, \\ \eta_3^{(4,r)} &= -\frac{1-\sqrt{r}-4r^{\frac{1}{4}}v_1}{4}, \end{aligned}$$

where u_1 *and* v_1 *are defined in Lemma 2.*
(b) If $p \equiv 3 \pmod 4$*, then* $m \equiv 0 \pmod 2$*.*
For the case of $m \equiv 0 \pmod 4$*, we have*

$$\begin{aligned} \eta_0^{(4,r)} &= -\frac{1+3\sqrt{r}}{4}, \\ \eta_j^{(4,r)} &= -\frac{1-\sqrt{r}}{4} \quad for \quad all \quad j \neq 0. \end{aligned}$$

For the case of $m \equiv 2 \pmod 4$, *we have*

$$\begin{aligned} \eta_0^{(4,r)} &= -\frac{1-3\sqrt{r}}{4}, \\ \eta_j^{(4,r)} &= -\frac{1+\sqrt{r}}{4} \quad for \quad all \quad j \neq 0. \end{aligned}$$

The following gives the complete weight enumerator of the code C_D with defining set D constructed from cyclotomy of order $N = 4$.

Theorem 9. *Let* $N = 4$, $r = p^m$ *and* $N | \frac{r-1}{p-1}$. *Let* $\#I = 3$ *and* $r - 1 = nN$. *Then the code* C_D *of* (1) *is a* $[3n, m]$ *linear code over* $\mathbb{F}_p$ *and its complete weight enumerator is given as follows.*
(a) If $p \equiv 1 \pmod 4$ *and* $m \equiv 0 \pmod 4$, *then*

$$\begin{aligned} CWE(C_D) &= w_0^{3n} + nw_0^{\frac{3n}{p} - \frac{p-1}{4p}(3-\sqrt{r}+2r^{\frac{1}{4}}u_1)} \prod_{\rho=1}^{p-1} w_\rho^{\frac{3n}{p} + \frac{1}{4p}(3-\sqrt{r}+2r^{\frac{1}{4}}u_1)} \\ &\quad + nw_0^{\frac{3n}{p} - \frac{p-1}{4p}(3-\sqrt{r}-2r^{\frac{1}{4}}u_1)} \prod_{\rho=1}^{p-1} w_\rho^{\frac{3n}{p} + \frac{1}{4p}(3-\sqrt{r}-2r^{\frac{1}{4}}u_1)} \\ &\quad + nw_0^{\frac{3n}{p} - \frac{p-1}{4p}(3+\sqrt{r}+4r^{\frac{1}{4}}v_1)} \prod_{\rho=1}^{p-1} w_\rho^{\frac{3n}{p} + \frac{1}{4p}(3+\sqrt{r}+4r^{\frac{1}{4}}v_1)} \\ &\quad + nw_0^{\frac{3n}{p} - \frac{p-1}{4p}(3+\sqrt{r}-4r^{\frac{1}{4}}v_1)} \prod_{\rho=1}^{p-1} w_\rho^{\frac{3n}{p} + \frac{1}{4p}(3+\sqrt{r}-4r^{\frac{1}{4}}v_1)}, \end{aligned}$$

where u_1 *and* v_1 *are defined in Lemma 2.*
(b) If $p \equiv 3 \pmod 4$ *and* $m \equiv 0 \pmod 4$, *then*

$$\begin{aligned} CWE(C_D) &= w_0^{3n} + 3nw_0^{\frac{3n}{p} - \frac{p-1}{4p}(3+\sqrt{r})} \prod_{\rho=1}^{p-1} w_\rho^{\frac{3n}{p} + \frac{1}{4p}(3+\sqrt{r})} \\ &\quad + nw_0^{\frac{3n}{p} - \frac{3(p-1)}{4p}(1-\sqrt{r})} \prod_{\rho=1}^{p-1} w_\rho^{\frac{3n}{p} + \frac{3}{4p}(1-\sqrt{r})}. \end{aligned}$$

(c) If $p \equiv 3 \pmod 4$ *and* $m \equiv 2 \pmod 4$, *then*

$$\begin{aligned} CWE(C_D) &= w_0^{3n} + 3nw_0^{\frac{3n}{p} - \frac{p-1}{4p}(3-\sqrt{r})} \prod_{\rho=1}^{p-1} w_\rho^{\frac{3n}{p} + \frac{1}{4p}(3-\sqrt{r})} \\ &\quad + nw_0^{\frac{3n}{p} - \frac{3(p-1)}{4p}(1+\sqrt{r})} \prod_{\rho=1}^{p-1} w_\rho^{\frac{3n}{p} + \frac{3}{4p}(1+\sqrt{r})}. \end{aligned}$$

Proof: From Lemma 8, we can deduce the following assertions.
(a) If $p \equiv 1 \pmod 4$ and $m \equiv 0 \pmod 4$, then

$$\begin{aligned} \eta_0^{(4,r)} + \eta_1^{(4,r)} + \eta_2^{(4,r)} &= -\frac{3+\sqrt{r}+4r^{\frac{1}{4}}v_1}{4}, \\ \eta_1^{(4,r)} + \eta_2^{(4,r)} + \eta_3^{(4,r)} &= -\frac{3-\sqrt{r}-2r^{\frac{1}{4}}u_1}{4}, \\ \eta_2^{(4,r)} + \eta_3^{(4,r)} + \eta_0^{(4,r)} &= -\frac{3+\sqrt{r}-4r^{\frac{1}{4}}v_1}{4}, \\ \eta_3^{(4,r)} + \eta_0^{(4,r)} + \eta_1^{(4,r)} &= -\frac{3-\sqrt{r}+2r^{\frac{1}{4}}u_1}{4}, \end{aligned}$$

where u_1 and v_1 are defined in Lemma 2.
(b) If $p \equiv 3 \pmod 4$ and $m \equiv 0 \pmod 4$, then

$$\begin{aligned}\eta_0^{(4,r)}+\eta_1^{(4,r)}+\eta_2^{(4,r)} &= \eta_2^{(4,r)}+\eta_3^{(4,r)}+\eta_0^{(4,r)}\\ &= \eta_3^{(4,r)}+\eta_0^{(4,r)}+\eta_1^{(4,r)}\\ &= -\frac{3+\sqrt{r}}{4},\\ \eta_1^{(4,r)}+\eta_2^{(4,r)}+\eta_3^{(4,r)} &= -\frac{3(1-\sqrt{r})}{4}.\end{aligned}$$

(c) If $p \equiv 3 \pmod 4$ and $m \equiv 2 \pmod 4$, then

$$\begin{aligned}\eta_0^{(4,r)}+\eta_1^{(4,r)}+\eta_2^{(4,r)} &= \eta_2^{(4,r)}+\eta_3^{(4,r)}+\eta_0^{(4,r)}\\ &= \eta_3^{(4,r)}+\eta_0^{(4,r)}+\eta_1^{(4,r)}\\ &= -\frac{3-\sqrt{r}}{4},\\ \eta_1^{(4,r)}+\eta_2^{(4,r)}+\eta_3^{(4,r)} &= -\frac{3(1+\sqrt{r})}{4}.\end{aligned}$$

The desired conclusions then follow from Equation (4). ■

Theorem 10. *Let $N=4$, $r=p^m$ and $N|\frac{r-1}{p-1}$. Let $\#I=2$ and $r-1=nN$. Then the code C_D of* (1) *is a $[2n,m]$ linear code over $\mathbb{F}_p$ and its complete weight enumerator is given as follows.*
(a) If $p \equiv 1 \pmod 4$ and $m \equiv 0 \pmod 4$, then

$$\begin{aligned}CWE(C_D) &= w_0^{2n}+2nw_0^{\frac{2n}{p}-\frac{p-1}{2p}(1+\sqrt{r})}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{1}{2p}(1+\sqrt{r})}\\ &\quad+2nw_0^{\frac{2n}{p}-\frac{p-1}{2p}(1-\sqrt{r})}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{1}{2p}(1-\sqrt{r})},\end{aligned}$$

for the case of $I=\{0,2\}$ and $I=\{1,3\}$, and for other cases,

$$\begin{aligned}CWE(C_D) &= w_0^{2n}+nw_0^{\frac{2n}{p}-\frac{p-1}{2p}(1+r^{\frac14}(u_1+2v_1))}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{1}{2p}(1+r^{\frac14}(u_1+2v_1))}\\ &\quad+nw_0^{\frac{2n}{p}-\frac{p-1}{2p}(1-r^{\frac14}(u_1+2v_1))}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{1}{2p}(1-r^{\frac14}(u_1+2v_1))}\\ &\quad+nw_0^{\frac{2n}{p}-\frac{p-1}{2p}(1+r^{\frac14}(u_1-2v_1))}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{1}{2p}(1+r^{\frac14}(u_1-2v_1))}\\ &\quad+nw_0^{\frac{2n}{p}-\frac{p-1}{2p}(1-r^{\frac14}(u_1-2v_1))}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{1}{2p}(1-r^{\frac14}(u_1-2v_1))},\end{aligned}$$

where u_1 and v_1 are defined in Lemma 2.
(b) If $p \equiv 3 \pmod 4$ and $m \equiv 0 \pmod 2$, then

$$\begin{aligned}CWE(C_D) &= w_0^{2n}+2nw_0^{\frac{2n}{p}-\frac{p-1}{2p}(1+\sqrt{r})}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{1}{2p}(1+\sqrt{r})}\\ &\quad+2nw_0^{\frac{2n}{p}-\frac{p-1}{2p}(1-\sqrt{r})}\prod_{\rho=1}^{p-1}w_\rho^{\frac{2n}{p}+\frac{1}{2p}(1-\sqrt{r})}.\end{aligned}$$

Proof: From Lemma 8, we can deduce the following assertions.
(a) If $p \equiv 1 \pmod 4$ and $m \equiv 0 \pmod 4$.

For the case of $I = \{0,2\}$ and $I = \{1,3\}$, we have

$$\begin{aligned} \eta_0^{(4,r)} + \eta_2^{(4,r)} &= -\tfrac{1+\sqrt{r}}{2} &= \eta_2^{(4,r)} + \eta_0^{(4,r)}, \\ \eta_1^{(4,r)} + \eta_3^{(4,r)} &= -\tfrac{1-\sqrt{r}}{2} &= \eta_3^{(4,r)} + \eta_1^{(4,r)}. \end{aligned}$$

And, for other cases, we have

$$\begin{aligned} \eta_0^{(4,r)} + \eta_1^{(4,r)} &= -\frac{1+r^{\frac{1}{4}}(u_1+2v_1)}{2}, \\ \eta_1^{(4,r)} + \eta_2^{(4,r)} &= -\frac{1-r^{\frac{1}{4}}(u_1-2v_1)}{2}, \\ \eta_2^{(4,r)} + \eta_3^{(4,r)} &= -\frac{1-r^{\frac{1}{4}}(u_1+2v_1)}{2}, \\ \eta_3^{(4,r)} + \eta_0^{(4,r)} &= -\frac{1+r^{\frac{1}{4}}(u_1-2v_1)}{2}. \end{aligned}$$

where u_1 and v_1 are defined in Lemma 2.
(b) If $p \equiv 3 \pmod 4$ and $m \equiv 0 \pmod 2$.
For the case of $m \equiv 0 \pmod 4$, we have

$$\begin{aligned} \eta_0^{(4,r)} + \eta_1^{(4,r)} &= -\tfrac{1+\sqrt{r}}{2} &= \eta_3^{(4,r)} + \eta_0^{(4,r)}, \\ \eta_1^{(4,r)} + \eta_2^{(4,r)} &= -\tfrac{1-\sqrt{r}}{2} &= \eta_2^{(4,r)} + \eta_3^{(4,r)}. \end{aligned}$$

For the case of $m \equiv 2 \pmod 4$, we have

$$\begin{aligned} \eta_0^{(4,r)} + \eta_1^{(4,r)} &= -\tfrac{1-\sqrt{r}}{2} &= \eta_3^{(4,r)} + \eta_0^{(4,r)}, \\ \eta_1^{(4,r)} + \eta_2^{(4,r)} &= -\tfrac{1+\sqrt{r}}{2} &= \eta_2^{(4,r)} + \eta_3^{(4,r)}. \end{aligned}$$

The desired conclusions then follow from Equation (4). ■

When $\#I = 1$, the result is straightforward from Lemma 8 and Equation (4).

Theorem 11. *Let $N = 4$, $r = p^m$ and $N | \frac{r-1}{p-1}$. Let $\#I = 1$ and $r - 1 = nN$. Then the code C_D of (1) is an $[n, m]$ cyclic code over $\mathbb{F}_p$ and its complete weight enumerator is given as follows.*
(a) If $p \equiv 1 \pmod 4$ and $m \equiv 0 \pmod 4$, then

$$\begin{aligned} CWE(C_D) &= w_0^n + n w_0^{\frac{n}{p} - \frac{p-1}{4p}(1+\sqrt{r}+2r^{\frac{1}{4}}u_1)} \prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{4p}(1+\sqrt{r}+2r^{\frac{1}{4}}u_1)} \\ &\quad + n w_0^{\frac{n}{p} - \frac{p-1}{4p}(1+\sqrt{r}-2r^{\frac{1}{4}}u_1)} \prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{4p}(1+\sqrt{r}-2r^{\frac{1}{4}}u_1)} \\ &\quad + n w_0^{\frac{n}{p} - \frac{p-1}{4p}(1-\sqrt{r}+4r^{\frac{1}{4}}v_1)} \prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{4p}(1-\sqrt{r}+4r^{\frac{1}{4}}v_1)} \\ &\quad + n w_0^{\frac{n}{p} - \frac{p-1}{4p}(1-\sqrt{r}-4r^{\frac{1}{4}}v_1)} \prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{4p}(1-\sqrt{r}-4r^{\frac{1}{4}}v_1)}, \end{aligned}$$

where u_1 and v_1 are defined in Lemma 2.
(b) If $p \equiv 3 \pmod 4$ and $m \equiv 0 \pmod 4$, then

$$\begin{aligned} CWE(C_D) &= w_0^n + 3n w_0^{\frac{n}{p} - \frac{p-1}{4p}(1-\sqrt{r})} \prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{4p}(1-\sqrt{r})} \\ &\quad + n w_0^{\frac{n}{p} - \frac{p-1}{4p}(1+3\sqrt{r})} \prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{4p}(1+3\sqrt{r})}. \end{aligned}$$

(c) If $p \equiv 3 \pmod 4$ *and* $m \equiv 2 \pmod 4$*, then*

$$\begin{aligned} CWE(C_D) &= w_0^n + 3nw_0^{\frac{n}{p}-\frac{p-1}{4p}(1+\sqrt{r})}\prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{4p}(1+\sqrt{r})} \\ &\quad + nw_0^{\frac{n}{p}-\frac{p-1}{4p}(1-3\sqrt{r})}\prod_{\rho=1}^{p-1} w_\rho^{\frac{n}{p}+\frac{1}{4p}(1-3\sqrt{r})}. \end{aligned}$$

Remark 12. We remark that the general strategy of Equation (3) is equivalent to the formula given in Theorem 3.1 of [26] for the special case of $D = C_0^{(N,r)}$. In other words,

$$\eta_k^{(N,r)} = \frac{1}{N}\sum_{i=0}^{N-1} G(\bar{\tau}^i)\tau^i(a), \tag{5}$$

where $a \in C_k^{(N,r)}$, $\tau = \chi^n$ and $\bar{\tau}$ be the conjugate character of τ. We shall show this in detail.

Recall that α is the primitive element of $\mathbb{F}_r$. Let $\eta_k^{(N,r)} = \sum_{x\in C_k^{(N,r)}} \psi(x)$, where $\psi(x) = \zeta_p^{\mathrm{Tr}(x)}$ is the canonical additive character over $\mathbb{F}_r$.

Define Gauss sum over $\mathbb{F}_r$ to be

$$G(\lambda) = \sum_{x\in\mathbb{F}_r^*} \lambda(x)\psi(x),$$

where λ is a multiplicative character of $\mathbb{F}_r$.

It is known that the set $\widehat{\mathbb{F}}_r^*$ of all the multiplicative characters of $\mathbb{F}_r^*$ forms a group generated by χ, i.e., $\widehat{\mathbb{F}}_r^* = \langle\chi\rangle$, where χ is a multiplicative character of order $r-1$. Then Gauss sums can be regarded as the Fourier coefficients in the Fourier expansion of the restriction of ψ to $\mathbb{F}_r^*$ in terms of the multiplicative characters of $\mathbb{F}_r$. That is

$$\psi(x) = \frac{1}{r-1}\sum_{\lambda\in\langle\chi\rangle} G(\bar{\lambda})\lambda(x), \quad \text{for} \quad x \in \mathbb{F}_r^*. \tag{6}$$

With above preparation, the left hand side of (6) is

$$\begin{aligned} LHS &= \sum_{x\in C_k^{(N,r)}} \psi(x) \\ &= \sum_{x\in C_k^{(N,r)}} \frac{1}{r-1}\sum_{\lambda\in\langle\chi\rangle} G(\bar{\lambda})\lambda(x) \\ &= \frac{1}{r-1}\sum_{\lambda\in\langle\chi\rangle} G(\bar{\lambda})\sum_{j=0}^{n-1}\lambda(\alpha^k\alpha^{Nj}) \\ &= \frac{1}{r-1}\sum_{\lambda\in\langle\chi\rangle} G(\bar{\lambda})\lambda(\alpha^k)\sum_{j=0}^{n-1}\lambda(\alpha^{Nj}) \\ &= \frac{n}{r-1}\sum_{\lambda\in\langle\chi^n\rangle} G(\bar{\lambda})\lambda(\alpha^k) \\ &= \frac{1}{N}\sum_{i=0}^{N-1} G(\bar{\chi}^{ni})\chi^{ni}(\alpha^k), \end{aligned}$$

where the fifth equal sign holds since

$$\sum_{j=0}^{n-1}\lambda(\alpha^{Nj}) = \begin{cases} n & \text{if} \quad \lambda^N = 1, \\ 0 & otherwise. \end{cases}$$

On the other hand the right hand side of (6) is

$$\begin{aligned}RHS &= \frac{1}{N}\sum_{i=0}^{N-1} G(\bar{\tau}^i)\tau^i(a)\\ &= \frac{1}{N}\sum_{i=0}^{N-1} G(\bar{\chi}^{ni})\chi^{ni}(\alpha^k\alpha^{Nj_0})\\ &= \frac{1}{N}\sum_{i=0}^{N-1} G(\bar{\chi}^{ni})\chi^{ni}(\alpha^k).\end{aligned}$$

Therefore, Equation (6) holds for all $x \neq 0$.

The results of case $p \equiv 2 \pmod 3$ in Theorem 5 and case $p \equiv 3 \pmod 4$ in Theorem 11 consistent with the result of Theorem 3.5 in [26] for semi-primitive case.

We also mention that the general strategy can be applied to the case of $N \in \{5, 6, 8, 12\}$, though we did not list them here, by ultilizing the Gaussian periods of order $N \in \{5, 6, 8, 12\}$ since the period polynomial $\Psi_{(N,r)}(X)$ and its factorization were determined explicitly for $N = 5$ in [40], and for $N \in \{6, 8, 12\}$ in [41], with quite complex expression.

Example 13. Let $(p, m) = (3, 4)$ and $N = 4$. Then $r = 81$ and $n = 20$. Suppose that α is a primitive element of $\mathbb{F}_{81}$.

(1) For the case of $D = C_1^{(4,r)} = \alpha\langle\alpha^4\rangle$.

By Theorem 11, the code C_D of (1) is a $[20, 4, 12]$ cyclic code over $\mathbb{F}_3$ with complete weight enumerator

$$CWE(C_D) = w_0^{20} + 60w_0^8w_1^6w_2^6 + 20w_0^2w_1^9w_2^9,$$

which is consistent with numerical computation by Magma.

This code is the best ternary cyclic code of length 20 and dimension 4 due to the tables given by Ding [2].

(2) For the case of $D = C_0^{(4,r)} \bigcup C_1^{(4,r)} = \langle\alpha^4\rangle \bigcup \alpha\langle\alpha^4\rangle$.

By Theorem 10, the code C_D of (1) is a $[40, 4, 24]$ linear code over $\mathbb{F}_3$ with complete weight enumerator

$$CWE(C_D) = w_0^{40} + 40w_0^{16}w_1^{12}w_2^{12} + 40w_0^{10}w_1^{15}w_2^{15},$$

which is consistent with numerical computation by Magma.

(3) For the case of $D = C_0^{(4,r)} \bigcup C_1^{(4,r)} \bigcup C_2^{(4,r)}$.

By Theorem 9, the code C_D of (1) is a $[60, 4, 36]$ linear code over $\mathbb{F}_3$ with complete weight enumerator

$$CWE(C_D) = w_0^{60} + 20w_0^{24}w_1^{18}w_2^{18} + 60w_0^{18}w_1^{21}w_2^{21},$$

which is consistent with numerical computation by Magma.

Example 14. Let $(p, m) = (5, 4)$ and $N = 4$. Then $r = 625$ and $n = 156$. Suppose that α is a primitive element of $\mathbb{F}_{625}$.

(1) For the case of $D = C_1^{(4,r)} = \alpha\langle\alpha^4\rangle$.

By Theorem 11, the code C_D of (1) is a $[156, 4, 112]$ cyclic code over $\mathbb{F}_5$ with complete weight enumerator

$$\begin{aligned}CWE(C_D) &= w_0^{156} + 156w_0^{44}(w_1w_2w_3w_4)^{28} + 156w_0^{32}(w_1w_2w_3w_4)^{31}\\ &\quad +156w_0^{28}(w_1w_2w_3w_4)^{32} + 156w_0^{20}(w_1w_2w_3w_4)^{34},\end{aligned}$$

which is consistent with numerical computation by Magma.

(2) For the case of $D = C_1^{(4,r)} \bigcup C_3^{(4,r)} = \alpha\langle\alpha^4\rangle \bigcup \alpha^3\langle\alpha^4\rangle$.

By Theorem 10, the code C_D of (1) is a $[312, 4, 240]$ linear code over $\mathbb{F}_5$ with complete weight enumerator

$$CWE(C_D) = w_0^{312} + 312w_0^{72}(w_1w_2w_3w_4)^{60} + 312w_0^{52}(w_1w_2w_3w_4)^{65},$$

which is consistent with numerical computation by Magma.

(3) For the case of $D = C_0^{(4,r)} \bigcup C_3^{(4,r)} = \langle\alpha^4\rangle \bigcup \alpha^3\langle\alpha^4\rangle$.

By Theorem 10, the code C_D of (1) is a $[312, 4, 236]$ linear code over $\mathbb{F}_5$ with complete weight enumerator

$$\begin{aligned}CWE(C_D) &= w_0^{312} + 156w_0^{76}(w_1w_2w_3w_4)^{59} + 156w_0^{64}(w_1w_2w_3w_4)^{62}\\ &\quad +156w_0^{60}(w_1w_2w_3w_4)^{63} + 156w_0^{48}(w_1w_2w_3w_4)^{66},\end{aligned}$$

which is consistent with numerical computation by Magma.

(4) For the case of $D = C_0^{(4,r)} \bigcup C_1^{(4,r)} \bigcup C_2^{(4,r)}$.

By Theorem 9, the code C_D of (1) is a $[468, 4, 364]$ linear code over $\mathbb{F}_5$ with complete weight enumerator

$$\begin{aligned} CWE(C_D) &= w_0^{468} + 156w_0^{104}(w_1w_2w_3w_4)^{91} + 156w_0^{96}(w_1w_2w_3w_4)^{93} \\ &\quad +156w_0^{92}(w_1w_2w_3w_4)^{94} + 156w_0^{80}(w_1w_2w_3w_4)^{97}, \end{aligned}$$

which is consistent with numerical computation by Magma.

IV. CONCLUDING REMARKS

In this paper, we proposed the complete weight enumerator of a family of linear code C_D with defining set D constructed from cyclotomy. The formulae for the general strategy and two special cases of $N = 3$ and $N = 4$ were presented by employing Gaussian periods. This indicates that the complete weight enumerator of C_D can be determined by the explicit Gaussian periods. As is well known that, the determination of Gaussian periods is quite complicated, so is the complete weight enumerator of C_D.

REFERENCES

[1] F. J. MacWilliams and N. J. A. Sloane, *The theory of error-correcting codes*. North-Holland Publishing, Amsterdam, 1977, vol. 16.

[2] C. Ding, *Codes from Difference Sets*. Singapore: World Scientific, 2014.

[3] I. F. Blake and K. Kith, "On the complete weight enumerator of Reed-Solomon codes," *SIAM J. Discret. Math.*, vol. 4, no. 2, pp. 164–171, Mar. 1991.

[4] D. Zheng, X. Wang, L. Yu, and H. Liu, "The weight enumerators of several classes of p-ary cyclic codes," *Discrete Mathematics*, vol. 338, no. 7, pp. 1264 – 1276, 2015.

[5] S. Yang, Z.-A. Yao, and C.-A. Zhao, "The weight enumerator of the duals of a class of cyclic codes with three zeros," *to appear in Applicable Algebra in Engineering, Communication and Computing*. [Online]. Available: http://link.springer.com/article/10.1007/s00200-015-0255-6

[6] C. Ding, Y. Liu, C. Ma, and L. Zeng, "The weight distributions of the duals of cyclic codes with two zeros," *IEEE Transactions on Information Theory*, vol. 57, no. 12, pp. 8000–8006, 2011.

[7] C. Ding and J. Yang, "Hamming weights in irreducible cyclic codes," *Discrete Mathematics*, vol. 313, no. 4, pp. 434–446, 2013.

[8] H. Q. Dinh, C. Li, and Q. Yue, "Recent progress on weight distributions of cyclic codes over finite fields," *Journal of Algebra Combinatorics Discrete Structures and Applications*, vol. 2, no. 1, pp. 39–63, 2015.

[9] K. Feng and J. Luo, "Weight distribution of some reducible cyclic codes," *Finite Fields and Their Applications*, vol. 14, no. 2, pp. 390–409, 2008.

[10] C. Li and Q. Yue, "Weight distributions of two classes of cyclic codes with respect to two distinct order elements," *IEEE Transactions on Information Theory*, vol. 60, no. 1, pp. 296–303, Jan 2014.

[11] J. Luo and K. Feng, "On the weight distributions of two classes of cyclic codes," *IEEE Transactions on Information Theory*, vol. 54, no. 12, pp. 5332–5344, Dec 2008.

[12] A. Sharma and G. K. Bakshi, "The weight distribution of some irreducible cyclic codes," *Finite Fields and Their Applications*, vol. 18, no. 1, pp. 144–159, 2012.

[13] G. Vega, "The weight distribution of an extended class of reducible cyclic codes," *IEEE Transactions on Information Theory*, vol. 58, no. 7, pp. 4862–4869, July 2012.

[14] B. Wang, C. Tang, Y. Qi, Y. Yang, and M. Xu, "The weight distributions of cyclic codes and elliptic curves," *IEEE Transactions on Information Theory*, vol. 58, no. 12, pp. 7253–7259, 2012.

[15] J. Yuan, C. Carlet, and C. Ding, "The weight distribution of a class of linear codes from perfect nonlinear functions," *IEEE Transactions on Information Theory*, vol. 52, no. 2, pp. 712–717, 2006.

[16] D. Zheng, X. Wang, X. Zeng, and L. Hu, "The weight distribution of a family of p-ary cyclic codes," *Designs, Codes and Cryptography*, vol. 75, no. 2, pp. 263–275, 2015.

[17] K. Kith, "Complete weight enumeration of Reed-Solomon codes," *Master's thesis, Department of Electrical and Computing Engineering, University of Waterloo, Waterloo, Ontario, Canada*, 1989.

[18] T. Helleseth and A. Kholosha, "Monomial and quadratic bent functions over the finite fields of odd characteristic," *IEEE Transactions on Information Theory*, vol. 52, no. 5, pp. 2018–2032, 2006.

[19] C. Ding, T. Helleseth, T. Klove, and X. Wang, "A generic construction of Cartesian authentication codes," *IEEE Transactions on Information Theory*, vol. 53, no. 6, pp. 2229–2235, 2007.

[20] C. Ding and X. Wang, "A coding theory construction of new systematic authentication codes," *Theoretical computer science*, vol. 330, no. 1, pp. 81–99, 2005.

[21] W. Chu, C. J. Colbourn, and P. Dukes, "On constant composition codes," *Discrete Applied Mathematics*, vol. 154, no. 6, pp. 912–929, 2006.

[22] C. Ding, "Optimal constant composition codes from zero-difference balanced functions," *IEEE Transactions on Information Theory*, vol. 54, no. 12, pp. 5766–5770, 2008.

[23] C. Ding and J. Yin, "A construction of optimal constant composition codes," *Designs, Codes and Cryptography*, vol. 40, no. 2, pp. 157–165, 2006.

[24] A. Kuzmin and A. Nechaev, "Complete weight enumerators of generalized Kerdock code and linear recursive codes over Galois ring," in *Workshop on coding and cryptography*, 1999, pp. 333–336.

[25] ——, "Complete weight enumerators of generalized Kerdock code and related linear codes over Galois ring," *Discrete applied mathematics*, vol. 111, no. 1, pp. 117–137, 2001.

[26] C. Li, Q. Yue, and F.-W. Fu, "Complete weight enumerators of some cyclic codes," *Designs, Codes and Cryptography*, pp. 1–21, 2015.

[27] C. Li, S. Bae, and J. Ahn, "Complete weight enumerators of linear codes and their applications," *preprint*.

[28] S. Bae, C. Li, and Q. Yue, "On the complete weight enumerators of some reducible cyclic codes," *Discrete Mathematics*, vol. 338, no. 12, pp. 2275 – 2287, 2015.

[29] S. Yang and Z. Yao, "Complete weight enumerators of some linear codes," 2015. [Online]. Available: http://arxiv.org/abs/1505.06326

[30] ——, "The complete weight enumerator of several cyclic codes," 2015. [Online]. Available: http://arxiv.org/abs/1505.05576

[31] C. Ding and H. Niederreiter, "Cyclotomic linear codes of order 3," *IEEE Transactions on Information Theory*, vol. 53, no. 6, pp. 2274–2277, June 2007.

[32] K. Ding and C. Ding, "Binary linear codes with three weights," *IEEE Communications Letters*, vol. 18, no. 11, pp. 1879–1882, Nov 2014.

[33] C. Ding, “Linear codes from some 2-designs,” *IEEE Transactions on Information Theory*, vol. 61, no. 6, pp. 3265–3275, June 2015.
[34] K. Ding and C. Ding, “A class of two-weight and three-weight codes and their applications in secret sharing,” *arxiv:1503.06512v1*.
[35] J. Yuan and C. Ding, “Secret sharing schemes from three classes of linear codes,” *IEEE Transactions on Information Theory*, vol. 52, no. 1, pp. 206–212, Jan 2006.
[36] C. Carlet, C. Ding, and J. Yuan, “Linear codes from perfect nonlinear mappings and their secret sharing schemes,” *IEEE Transactions on Information Theory*, vol. 51, no. 6, pp. 2089–2102, 2005.
[37] J. Luo and T. Helleseth, “Constant composition codes as subcodes of cyclic codes,” *IEEE Transactions on Information Theory*, vol. 57, no. 11, pp. 7482–7488, Nov 2011.
[38] O. Milenkovic and N. Kashyap, “On the design of codes for dna computing,” in *Coding and Cryptography*, ser. Lecture Notes in Computer Science. Springer Berlin Heidelberg, 2006, vol. 3969, pp. 100–119.
[39] G. Myerson, “Period polynomials and Gauss sums for finite fields,” *Acta Arithmetica*, vol. 39, no. 3, pp. 251–264, 1981.
[40] A. Hoshi, “Explicit lifts of quintic Jacobi sums and period polynomials for $\mathtt{F_q}$,” *Proceedings of the Japan Academy, Series A, Mathematical Sciences*, vol. 82, no. 7, pp. 87–92, 2006.
[41] S. J. Gurak, “Period polynomials for $\mathbb{F}_q$ of fixed small degree,” *CRM Proc. and Lect. Notes*, vol. 36, pp. 127–145, 2004.